



(A) A presurgical angiogram of the left common carotid artery showed tapered occlusion of the internal carotid artery (arrow).
 (B) The internal carotid artery was shown on angiogram to be patent after mechanical thrombectomy.

for management of AIS among pediatric patients (younger than 18 years of age) are based on data extrapolated from adult data. The use of intravenous thrombolytic agents such as tPA is considered experimental given the lack of clinical trials and prospective studies of its use in the pediatric population. However, results of the Thrombolysis in Pediatric Stroke (TIPS) trial,⁴ a 5-year prospective trial examining the use of intravenous tPA in children aged 2 to 17 years, will be crucial for creating evidence-based recommendations for treating pediatric patients with AIS. ■

REFERENCES

1. Lyle CA, Bernard TJ, Goldenberg NA. Childhood arterial ischemic stroke: a review of etiologies, antithrombotic treatments, prognostic factors, and priorities for future research. *Semin Thromb Hemost.* 2011;37(7):786-793.
2. Mackay MT, Wiznitzer M, Benedict SL, Lee KJ, deVeber GA, Ganesan V; International Pediatric Stroke Study Group. Arterial ischemic stroke risk factors: the International Pediatric Stroke Study. *Ann Neurol.* 2011;69(1):130-140.
3. Gdynia H-J, Kühnlein P, Ludolph AC, Huber R. Connective tissue disorders in dissections of the carotid or vertebral arteries. *J Clin Neurosci.* 2008;15(5):489-494.
4. Ellis MJ, Amlie-Lefond C, Orbach DB. Endovascular therapy in children with acute ischemic stroke: review and recommendations. *Neurology.* 2012;79(13 suppl 1):S158-S164.

Benign Primary Milia

Alexander K. C. Leung, MD
 University of Calgary

Benjamin Barankin, MD
 Toronto Dermatology Centre

A 17-year-old presented with multiple white papules on her face that had been first noted approximately 4 years ago. These asymptomatic papules had appeared gradually over 18 months' time. Her medical history was otherwise unremarkable. In particular, there was no

history of trauma, and she was not on any topical or oral medication. A diagnosis of benign primary milia of children was made.

Milia are small (generally less than 3 mm), white, benign, dome-shaped, superficial keratinous cysts. Histologically, they appear as small infundibular cysts that are lined with stratified squamous epithelium with a granular cell layer.¹ The cyst contains laminated layers of keratin. Milia may arise spontaneously without known cause (primary milia) or secondary to various processes (secondary milia). Primary milia may be congenital or have onset later in life.

Congenital primary milia are present in approximately 40% of newborn infants, with no sex predilection. The condition is less common and is delayed in onset in premature infants.¹ While congenital primary milia favor the nose, benign primary milia of children and adults favor the eyelids. Other unusual sites of involvement include the areola, nasal crease, glans penis, and vulva.¹ The presence of primary milia may be a feature of Rombo syndrome, Brooke-Spiegler syndrome, Bazex-Dupré-Christol syndrome, Basan syndrome, orofaciocigital syndrome type 1, basal cell nevus syndrome, and Nicolau-Balus syndrome.¹



The differential diagnosis of primary milia includes sebaceous hyperplasia, syringomas, molluscum contagiosum, lichen nitidus, closed comedones, cutaneous myxomas, and calcinosis cutis.¹

Milia en plaque and multiple eruptive milia are considered variants of milia.^{2,3} Milia en plaque is characterized by a plaque formed by a confluence of closely grouped, tiny, skin-colored to white papules, often with an erythematous base.³ The condition is most common in middle-aged individuals, with a female predominance and no predilection for race.³ The lesions usually are asymptomatic. The postauricular area is most commonly affected. Other sites of involvement include the periocular area, supraclavicular area, submandibular area, cheeks, nasal bridge, trunk, and lower limbs.³ Multiple eruptive milia are characterized by a sudden onset of multiple milia over a period of weeks to months, mainly on the head, neck, and trunk.² The lesions usually are asymptomatic.²

Secondary milia may occur in association with disease (eg, epidermolysis bullosa, porphyria cutanea tarda), medication (eg, oral cyclosporine, penicillamine, 5-fluorouracil, topical corticosteroids), or trauma (dermabrasion, chemical peels, burns, skin grafts, tattoos).⁴

Primary congenital milia often exfoliate and resolve spontaneously in the first few weeks to months of life, thus requiring no treatment. Benign primary milia of children and adults tend to persist longer. The lesions of milia en plaque and multiple eruptive milia usually persist without treatment.³ For secondary milia, the underlying cause has to be removed if possible. For persistent lesions, treatment options include simple mechanical evacuation or extraction, topical retinoids, electrocautery, electrodesiccation, cryosurgery, or carbon dioxide laser vaporization.⁵ ■

REFERENCES

1. Berk DR, Bayliss SJ. Milia: a review and classification. *J Am Acad Dermatol*. 2008;59(6):1050-1063.
2. Feet N, Nevas J. Multiple eruptive milia. *Cutis*. 2013;91(4):191-192.
3. Zhang R-Z, Zhu W-Y. Bilateral milia en plaque in a 6-year-old Chinese boy. *Pediatr Dermatol*. 2012; 29(4):504-506.
4. Miller LM, Schwartz JT, Cho S. Milia: a unique reaction tattoos. *Cutis*. 2011;87(4):195-196.
5. Del Pozo J, Castiñeiras I, Fernández-Jorge B. Variants of milia successfully treated with CO₂ laser vaporization. *J Cosmet Laser Ther*. 2010;12(4):191-194.

Professional Opportunities

Pediatric practice for sale in Southwest Florida – Sarasota/Bradenton area.

Contact Miguel Alvarez –
mickiesdiscus@outlook.com

For information on placing a classified advertisement, contact Joseph Miller at jmiller@hmpcommunications.com or Mackenzie Berry at mberry@hmpcommunications.com

Consultant FOR PEDIATRICIANS

ADVERTISER INDEX

April 2014

Shire

Vyvance Cover 2

American Lifeline

Florajen4Kids 149

Johnson & Johnson

Desitin 151

McNeil

Children's Zyrtec 153

Merck

PedvaxHIB Insert

Merck

RotaTeq 162

Galderma Laboratories

Epiduo Insert

Salix Pharmaceuticals

Deflux Insert

Summers Laboratories

Triple Paste 185

Mead Johnson Nutrition

Enfamil Reguline Cover 3

Lansinoh

Breastfeeding and Baby Products Cover 4