

4256 Bathurst St., Suite 400 Toronto, ON, M3H 5Y8 Tel.: (416) 633-0001

Fax: (416) 633-0002

www.torontodermatologycentre.com info@torontodermatologycentre.com

WOUND CARE INSTRUCTIONS AFTER SKIN BIOPSY OR SURGERY

- Keep the area dry for at least 24 hours, although 48 hours is better. After 48 hours you **MUST** wash the area daily with soap and water (if you have a beige outer bandage remove it at this time without removing the steri-strips underneath). Avoid immersing the incision in ocean or swimming pool or jacuzzi water for 2 weeks.
- If you have Steri-Strips, allow for them to fall off naturally. If they fall off in less than 10 days start applying antibiotic ointment (e.g. Fucidin, Bactroban, Polysporin) or Vaseline ointment 2-4 times per day till 10 days post-procedure.
- You should also cover the treated area if you will be in dirty environment or if there is a chance the wound will be injured or traumatized (e.g. at work, under clothing that is rubbing, playing sports, etc).
- If there is excessive bleeding, apply pressure for 20 minutes uninterrupted. If bleeding fails to stop, contact the Toronto Dermatology Centre. If it is after hours or weekend, go to nearest walk-in clinic, GP or Emergency Room.
- Do not pick off scabs new ones will form if the old ones are removed too soon and it increased the possibility of scarring. Don't scrub or rub the wound until it heals.
- Activity: avoid any activity or movement that may cause the wound to stretch or place tension on the wound for 2-3 weeks. This will help prevent the wound from opening and will result in a finer, less visible scar that is less prone to widening over time. Sutured scar lines do not reach their full strength for 6 weeks. Once the stitches are removed the wound is no longer supported so to avoid stretching the scar, avoid movements which stretch across the suture line. In a small percentage of cases the suture site can open. This is no cause for concern but please start applying antibiotic ointment 2-3 times per day until the wound closes. This can take 1-2 weeks. Please note we will not re-stich a procedure site that has opened as that would increase the chance of an infection occurring.
- Make-up/hair colour: don't apply make-up to your wound until the skin has healed over and there is no scab (e.g. no sooner than 2 weeks). Don't apply make-up over stitches. If the procedure site is on your scalp, avoid colouring your hair until the area has healed.
- If you notice signs of infection which include: redness extending well beyond the surgery site, swelling, increasing pain, fever or sweats, odour, or pus draining from the site, please contact your family doctor or the nurse at the Toronto Dermatology Centre at 416-633-0001 x 3.
- Stitch removal: if you have had non-dissolvable stitches, they should be generally removed after: 5-7 days on the face or genitals; 7-10 days on the scalp; 10-14 days on the rest of the body
- o Make sure to book an appointment for stitch removal with our nurse at the appropriate time.
- o Note that dissolvable stitches can take anywhere from 2-4+ weeks to completely dissolve on their own.
- If concerned about scar appearance, we recommend Scar Esthetique Scar Crème or SkinMedica's Scar Recovery Gel. It will improve the scar appearance by softening and lightening it. Please note you must wait a minimum of 3 weeks post procedure to start using any scar minimizing product. Ask our receptionists or medical estheticians if interested.
- Pathology reports can take 2 weeks and you are **only** contacted if results indicate that further follow-up is required.
- All these instructions can be found on our website at www.torontodermatologycentre.com